



PATIENT INFORMATION

Patient Name: _____
 Address: _____
 Servicing Address: _____
 Phone Number: _____ Date of Birth: (MM/DD/YYYY) _____
 E-Mail: _____ Gender: Male Female Unspecified
 Preferred Method of Communication: Call Text Email

PRESCRIBER INFORMATION

Prescriber Name: _____
 Address: _____
 Primary Contact: _____ NPI: _____
 Contact Phone: _____ Tax ID: _____
 Contact Email: _____ MCD #: _____
 ICD-10 Diagnosis: _____

FACILITY INFORMATION

Facility Name: _____ Primary Contact Name: _____
 Contact Email: _____ Contact Phone: _____
 Address: _____
 White Bag Delivery Address (in-facility infusions only): _____
 NPI #: _____ Tax ID #: _____ MCD #: _____

SITE OF CARE

- Home Office Independent Clinic Inpatient
 Off-Campus Outpatient Hospital On-Campus Outpatient Hospital Ambulatory Infusion
 Billing Codes: Home (12), Office (11), Independent Clinic (49), Inpatient (21), Off-Campus Outpatient Hospital (19), On-Campus Hospital (22), Ambulatory Infusion (24)

TREATMENT REGIMEN

Expected Cisplatin Start of Care Date

Cisplatin cut off time 2PM local time

Sample Dose Calculation: Actual Patient Body Weight: 24.9 kg Body Surface Area (BSA): 1.09 m² PEDMARK Dose: 20 g/m²
Dose based on the patient's body weight; see chart below.

Calculation: Multiply the patient's BSA by the PEDMARK Dose: **1.09 m² x 20 g/m² = 21.8 g total PEDMARK to be administered.**

PEDMARK DOSE

Weight: _____ kg Height: _____ cm BSA: _____ m²
 10g/m² (weight less than 5kg)
 15g/m² (weight of 5 to 10kg)
 20g/m² (weight greater than 10kg)

PRESCRIPTION

Infuse PEDMARK _____ grams IV over 15 minutes OR _____ (30-60) minutes to be administered
 6 hours after completion of cisplatin infusions that are 1 to 6 hours in duration.
 Dispense: Quantity sufficient of Pedmark (SDV) for each dose Refills: 12 months
 Cisplatin regimen: _____ (ex: 100mg/m² Q21D x2)

PEDMARK INFUSION ORDERS

PROVIDER SIGNATURE

Dispense as Written Signature _____ Date of Signature _____

Cisplatin infusion should be scheduled and prioritized to ensure completion by 2 PM local time to support same-day PEDMARK administration.

IMPORTANT NOTICE: This facsimile transmission is intended to be delivered only to the named addressee and may contain material that is confidential, privileged, proprietary or exempt from disclosure under applicable law. If it is received by anyone other than the named addressee, the recipient should immediately notify the sender at the address and telephone number set forth herein and obtain instructions as to disposal of the transmitted material. In no event should such material be read or retained by anyone other than the named addressee, except by the express authority of the sender to the named addressee.

REQUIRED DOCUMENTATION

CLINICAL DOCUMENTATION:

- RELEVANT CLINICAL NOTES OR TREATMENT PLAN
- DOCUMENTATION OF CURRENT CHEMOTHERAPY REGIMEN AND NEED FOR PEDMARK
- STATEMENT INDICATING PEDMARK IS NECESSARY TO REDUCE RISK OF CISPLATIN OTOTOXICITY
- MOST RECENT LAB RESULTS
- ENT/AUDIOLOGY NOTES AND RESULTS (if applicable)
- CENTRAL VENOUS ACCESS PLACEMENT/VERIFICATION DOCUMENT

INTAKE DOCUMENTATION:

- PEDMARK ENROLLMENT FORM / INFUSION ORDER
- COPY OF FRONT AND BACK OF INSURANCE CARD
- PATIENT AUTHORIZATION FORM (if available)
- FACE SHEET (Patient demographics, insurance details, contact information)

Complete This Section If Home Infusion
FENNEC HEARS ENROLLMENT FORM

ACCESS TYPE

(Also include Peripheral IV for Port Malfunction)

VENOUS ACCESS

- Peripheral IV
 - Peripheral IV may remain in place for multi-day home infusion therapy
 - Remove PIV at the end of the Pedmark infusion _____
provider initial
- Port _____ G _____ Inch Needle
 - De-access on last day of multi-day home infusion therapy
 - De-access after every Pedmark infusion
 - De-access per patient request
- PICC

VERIFICATION OF PLACEMENT (CENTRAL VENOUS ACCESS ONLY)

- Number of Lumens: 1 2 3
- Method of catheter placement verification: _____
(i.e. Radiological report/pst-insertion, chest x-ray, ECG verification for PICC, Ultrasound, Fluoroscopy)
 - Above central venous catheter successfully placed on _____ and viable
date of placement
 - Placement verified by _____
verified by (provider name and title)
- If no documentation available, has the catheter previously been used for infusion without complication?* _____
provider initial

FLUSH PROTOCOL

(If Heparin Flush is not selected OK to Saline lock port)

SALINE FLUSH

- 0.9% Saline Flush:** **Refills: 12 months**
Dispense: Quantity sufficient for each infusion
Flush catheter with 5-10mL for patency/SASH protocol

CATHETER FLUSH ORDERS

- Heparin Flush:** **Refills: 12 months**
Dispense: Quantity sufficient for each infusion
Flush catheter with 5mL of Heparin 100 units/mL per SASH protocol.

PEDMARK PRE-MEDICATION

Dispense: Quantity Sufficient of SDV or Tablet/Capsule for Each Dose | Refills 12 Months

Prescribing physician will use their independent professional judgement for each individual patient's treatment recommendations

At-home Infusions	Check to Include	Pre-Medication	Provider Selected Dose (mg)	Admin Route (IV or PO)	Provider Notes	Administrative Notes
Night Before Pedmark:	<input type="checkbox"/>	Olanzapine (Zyprexa) 5mg-10mg		<input type="checkbox"/> PO		Not shipped for home infusion
30-60 mins Before Pedmark:	<input type="checkbox"/>	Lorazepam (Ativan) 0.5mg-1.0mg		<input type="checkbox"/> PO		Not shipped for home infusion
	<input type="checkbox"/>	Famotidine (Pepcid) 20mg		<input type="checkbox"/> PO <input type="checkbox"/> IV		
	<input type="checkbox"/>	LMX 4 or (EMLA) Lidocaine 2.5%/Prilocaine 2.5% cream				Apply topically 1 hour prior to starting IV or accessing port QTY: 1
	<input type="checkbox"/>	Diphenhydramine (Benadryl) 25mg		<input type="checkbox"/> PO <input type="checkbox"/> IV		
	<input type="checkbox"/>	Dexamethasone (Decadron) 10mg-20mg		<input type="checkbox"/> PO <input type="checkbox"/> IV		20 mg max daily dose
	<input type="checkbox"/>	Ondansetron (Zofran) 8mg		<input checked="" type="checkbox"/> IV		24 mg max daily dose
	<input type="checkbox"/>	Metoclopramide (Reglan) 10mg		<input type="checkbox"/> PO <input type="checkbox"/> IV		
During Pedmark, if needed:	<input type="checkbox"/>	Prochlorperazine (Compazine) 5mg-10mg		<input checked="" type="checkbox"/> IV		
	<input type="checkbox"/>	List:		<input type="checkbox"/> PO <input type="checkbox"/> IV		
Other Medication(s):	<input type="checkbox"/>	Diphenhydramine (Benadryl) 12.5mg-25mg		<input checked="" type="checkbox"/> IV		E.g. infusion related reaction, breakthrough
	<input type="checkbox"/>	List:		<input type="checkbox"/> PO <input type="checkbox"/> IV		
	<input type="checkbox"/>	List:		<input type="checkbox"/> PO <input type="checkbox"/> IV		
	<input type="checkbox"/>	List:		<input type="checkbox"/> PO <input type="checkbox"/> IV		

SKILLED NURSING VISIT

Expected Cisplatin Infusion Start Time _____ Expected Cisplatin Infusion End Time _____

As needed for IV access, administration, and proper clinical monitoring. Administration procedures to be followed per pharmacy protocol.

- Post Infusion: Flush IV with 15 mL 0.9% Sodium Chloride Injection, USP at final rate of drug infusion.
- Vital Signs: At baseline and at every _____ minutes during infusion, at completion of post-infusion flush and 30 minutes after completion of post-infusion flush.
- Supplies: Provide infusion pump, IV Pole, back-up peripheral IV kit and all necessary infusion supplies, as necessary, if home health.

EMERGENCY MEDICATIONS

INSTRUCTIONS DURING REACTION

Dispense: 30 days Refills: 12 months

- 1) STOP PEDMARK
- 2) START OR ADMINISTER EMERGENCY MEDICATIONS
 - Diphenhydramine 25 mg (>30 kg) or 1.25 mg/kg (≤30 kg) IV or IM; repeat x 1 in 15 min PRN
 - Epinephrine 0.3 mg (> 30 kg), 0.15 mg (15 to 30 kg), or 0.01 mg/kg (< 15 kg) SQ or IM x 1; repeat x 1 in 5 to 15 min PRN
 - Corticosteroid - please specify (i.e. hydrocortisone, methylprednisolone): _____ Dose _____
 - 0.9% Sodium Chloride 500 mL (>30 kg) or 250 mL (<30 kg) IV at KVO rate PRN anaphylaxis: _____
 - 0.9% Saline _____ mL - administer at _____ mL/hr once PEDMARK infusion stops: _____
 - Other Emergency Medication(s): _____

3) CALL PHYSICIAN

- RESUME PEDMARK INFUSION AT SAME STARTING RATE ONCE PATIENT STABLE

PEDMARK PRE-MEDICATION AND INFUSION REACTION MANAGEMENT ORDERS

Dispense as Written Signature _____ Date of Signature _____ Product Substitution Permitted Signature _____ Date of Signature _____

IMPORTANT NOTICE: This facsimile transmission is intended to be delivered only to the named addressee and may contain material that is confidential, privileged, proprietary or exempt from disclosure under applicable law. If it is received by anyone other than the named addressee, the recipient should immediately notify the sender at the address and telephone number set forth herein and obtain instructions as to disposal of the transmitted material. In no event should such material be read or retained by anyone other than the named addressee, except by the express authority of the sender to the named addressee.

[Click Here for Prescribing Information](#)